

Determinant Calculations with Random Walk Worldline Loops

Michael G. Schmidt^{a 1} and Ion-Olimpiu Stamatescu^{a,b 2}

^a*Institut für Theoretische Physik, Philosophenweg 16, D-69120 Heidelberg, Germany*

^b*Forsch. der Ev. Studiengemeinschaft, Schmeilweg 5, D-69118, Heidelberg, Germany*

Abstract

We propose to calculate bosonic and fermionic determinants with some general field background, and the corresponding 1-loop effective actions by evaluating random walk worldline loops generated statistically on the lattice. This is illustrated by some numerical calculations for simple gauge field backgrounds and then discussed for the general case.

¹m.g.schmidt@thphys.uni-heidelberg.de

²stamates@thphys.uni-heidelberg.de

1 Introduction

The discussion of bosonic and fermionic determinants in some background and of the related 1-loop effective action is most transparent in the worldline/relativistic particle formalism [1]-[3]. The Euler-Heisenberg effective action of a constant electromagnetic field [4], for instance, can be presented very elegantly in this way [3]. Also for general gauge field backgrounds this formulation is very helpful allowing for an economical inverse mass (derivative) expansion to high order [5]. For background fields like an instanton, sphaleron or a bounce solution used in tunneling problems the evaluation of a derivative expansion [6] and alternatively the approach based on solutions for the eigenvalue spectrum [7] are rather sophisticated. The nonabelian gauge field groundstate background in 3- and 4-dimensional QCD and in the hot electroweak theory is even less accessible to an analytical treatment. For instance, the discussion of a nonperturbative magnetic and tachyonic mass in 3- and 4-dimensional SU(2) theory is very difficult even if one uses simplified assumptions about the gauge field vacuum [8]. Finally, we should notice that 2-loop effective actions are also accessible [9, 3, 10]

In this paper we consider the numerical evaluation of quantum field theoretical determinants in the worldline formalism using lattice methods. We consider the bosonic determinant in a gauge field background

$$\begin{aligned}
\Gamma(A) &= -\log \det(\mathcal{D}^2 + m^2) = -\text{tr} \log(\mathcal{D}^2 + m^2) \\
&= \int_0^\infty \frac{dT}{T} \int d^D x \int_{x(0)=x(T)=x} [Dx] \text{tr}_P \exp \left\{ - \int_0^T d\tau \left(\frac{\dot{x}^2}{4} + i g \mathbf{A}_\mu \dot{x}_\mu + m^2 \right) \right\} \\
&= \int d^D x \int_0^\infty \frac{dT}{T} e^{-m^2 T} \int_{x(0)=x(T)=x} [Dx] \exp \left\{ - \int_0^T d\tau \frac{\dot{x}^2}{4} \right\} \\
&\quad \times \text{tr}_P \exp \left\{ - i g \oint \mathbf{A}_\mu dx_\mu \right\}, \tag{1}
\end{aligned}$$

where the worldline sum (here already in the Euclidean formulation adequate for the lattice analysis) is over all closed paths and tr_P indicates path ordering. In proceeding to the lattice formulation of the problem it is natural to consider the loops on the same (usually hypercubic) lattice, on which the general gauge field configurations are generated. This appears to be difficult if one wants to enforce the $\exp\{-\int_0^T d\tau \frac{\dot{x}^2}{4}\}$ weight for randomly chosen paths, as proposed in an interesting recent paper [11]. However, as is well known, random walk paths automatically implement the above integral weight of the free particle propagation on a worldline [12]. The expression (1) can thus be discretized on a D-dimensional lattice Λ_D as

$$\Gamma^{discr}(A) = \sum_{x \in \Lambda_D} \mathcal{L}(A, x); \quad \mathcal{L}(A, x) = \sum_L \frac{1}{L} e^{-m^2 \frac{L}{2D}} \left(\frac{1}{2D} \right)^L \sum_{\{\omega_L(x)\}} \text{tr} \left(\prod_{l \in \omega_L(x)} U_l \right), \tag{2}$$

where dimensionfull quantities are understood as given in units of the lattice scale a according to their naive dimension and we have also discretized T using the τ intervals

$$\Delta\tau = \frac{T}{L} = \frac{a^2}{2D} \tag{3}$$

Here $\{\omega_L(x)\}$ is the set of all closed lattice paths with length $|\omega_L| = L$ obtained by random walk (RW) starting and ending at x , and we have written down explicitly the random walk “measure” $(2D)^{-L}$ (to be implemented by the actual RW procedure). The mass dependence $e^{-m^2 T}$ could be also obtained by a random walk weight but we prefer to keep this dependence explicit. U_l are link variables.

On the lattice the expansion according to the loop lengths (2) is of course equivalent to the usual hopping parameter expansion for the logarithm of the determinant, both for bosons and for fermions (see, e.g., [12], [13], [14]). This can be seen explicitly by rewriting (e.g., in the bosonic case)

$$\kappa^L = (m^2 + 2D)^{-L} \underset{m^2 \ll 2D}{\simeq} (2D)^{-L} e^{-m^2 \frac{L}{2D}} \quad (4)$$

(and similarly for fermions). The number of different loops of length L increases exponentially with the length, therefore one usually must cut the hopping parameter expansion at a rather small L . We suggest here to use a statistical procedure for sampling loops in a wide interval of lengths, instead of summing over all possible loops up to a given order (see also [15]). In our approach we use the random walk representation (2) to construct a (*sub-*)ensemble of loops of various lengths and evaluate the determinant with help of this sub-ensemble. By construction, the loops appear with the correct probabilities, and since this is a statistical procedure we may expect to have only statistical errors. Since loops of length L are typically relevant at the scale $\sqrt{L}$, arbitrarily cutting the loop expansion at some limited L may completely miss physical effects at scales larger than $\sqrt{L}$. In our method we trade these systematic errors for statistical ones such that we may hope to reproduce also such effects, albeit within statistical uncertainties. Notice that, on the one hand, we cannot use too rough lattices since this introduces discretization errors. On the other hand, the contribution of large loops is damped even at small mass by a factor $\propto L^{-D/2}$, see (8) next section. This suggests that, depending on the effects we want to observe, there is an optimal range of loops to be taken into account. In practice we use loops of length up to $L \simeq 200$ on lattices of 32^D . For the cases studied here these ensembles appeared adequate.

Chapter 2 describes the procedure how to generate closed loops and checks that the case without interaction on our loop ensembles converges to the well-known continuum (large L) result. In expression (2) we can insert some (quasi) classical background gauge field A_μ discretized by lattice connections U . This we will do for a constant magnetic field in chapter 3. This case allows comparison with the well-known Euler-Heisenberg action in its worldline formulation. We also treat there the case of constant $SU(2)$ fields in two perpendicular 2-planes in four dimensions which has a topological meaning [16]. The case of a constant magnetic field in a half space treated in ref. [11], which introduces a correlation scale (penetration length) is also studied.

For a quantum gauge field vacuum the action $\Gamma^{discr}(A)$ has to be inserted into a gauge field path integral discretized on the lattice to obtain the (quenched) vacuum energy

$$V(\beta) = \langle \Gamma \rangle = \sum_L \frac{1}{L} e^{-m^2 \frac{L}{2D}} \left(\frac{1}{2D} \right)^L \sum_{\{\omega_L\}} \frac{1}{Z_{YM}} \int [DU] e^{-\beta S_{YM}(\{U\})} \text{tr} \left(\prod_{l \in \omega_L} U_l \right) \quad (5)$$

with the Yang-Mills discretized action S_{YM} and partition function Z_{YM} . In the case of a further field ϕ in the background coupling to the loop a term $V''(\phi(x))$ has to be introduced in the discretized worldline action and further weights the sum over random paths. These 1-loop actions in a general background are our real goal to be accomplished in future work. Chapter 4 presents a final discussion and an outlook to the case of spin in the loop, in particular to fermionic determinants.

2 The lattice loop ensemble

As remarked in the introduction we use random walk to produce a loop ensemble thermalized with the “Boltzmann factor”

$$e^{-\frac{1}{4} \sum_{i=0}^{L-1} \sum_{\mu=1}^D (n_{\mu}(i+1) - n_{\mu}(i))^2 |_{n_{\mu}(0)=n_{\mu}(L)}}, \quad (6)$$

see (2), which is the discretized form of the corresponding factor in (1). In reconstructing the whole path integral we only need to weight the RW loop contributions $\text{tr} \left(\prod_{l \in \omega_L} U_l \right)$ with

$$L^{-1} e^{-m^2 \frac{L}{2D}} \quad (7)$$

since the factor $(2D)^{-L}$ is implicit in the RW generation. From now on we denote by $\{\omega\}$, $\{\omega_L\}$ the corresponding ensemble of loops *as it is produced* by the random walk procedure, that is, with the frequencies with which the loops are generated in the actual procedure.

In generating the loops we specify a maximal number of trials $\mathcal{N}$ (e.g., 2000000 for the 32^3 lattice) for each given number of steps (between 4 and 180). We first generate an $\{\omega(x_{\mu} = 1)\}$ ensemble of loops by starting the random walk at the point $x_{\mu} = 1$ of the periodic lattice and collecting the loops which after the prefixed number of steps have returned to $x_{\mu} = 1$, independently on possible self-crossings or retracing. We define $\mathcal{L}(A, 1)$ using this $\{\omega(x_{\mu} = 1)\}$ ensemble (for the application in sect. 3.2 we first shift the approximate “center of mass” of the loops to $x_{\mu} = 1$). For any given x we can calculate $\mathcal{L}(A, x)$ in (2) by accordingly shifting the loops of the $\{\omega(x_{\mu} = 1)\}$ ensemble as a whole. On the other hand, the full loop ensemble, that is, the x -summation $\sum_{x \in \Lambda_D}$ in (2), is approximated by considering sufficiently many random translations of each of the loops of the $\{\omega(x_{\mu} = 1)\}$ ensemble (for homogeneous field configurations this is, of course, redundant). That is, loop and x “summations” are done simultaneously. This construction, i.e. generation of loops at $x = 1$ and stochastic summation over x by random translations, automatically reproduces (statistically) the correct loop multiplicities in the hopping parameter expansion, by which a certain loop of length L appears L times, unless it represents r windings of a loop of length L' , $L = r L'$, in which case it only appears L' times [12], [14]. To improve the loop ensemble we can implement random rotations of the loops (before translations). Since the loop evaluation is vectorized over rotations and translations, these are reasonably cheap. In the following we work in $D = 3$ and $D = 4$ and use an $\{\omega(x_{\mu} = 1)\}$ ensemble of up to 600000 (250000) loops on lattices of $N_x \times N_y \times N_z (\times N_t)$ with $N_{\mu} = 12, 16, 24$ and 32 , and periodic boundary conditions.

For a free theory ($U_l = 1$ above) the contribution of the RW-loops of length L is proportional to their number, n_L (which counts the different loops with their multiplicities). From (1) we have:

$$n_L \simeq T^{-D/2} \propto L^{-D/2} \quad (8)$$




Figure 1: Number of closed loops of length L , n_L generated by random walk. We show the frequencies ν_L (9) vs L in $D = 3$ (diamonds) and $D = 4$ dimensions (crosses). The full lines represent the corresponding continuum results $L^{-D/2}$.

for large L – this well known relation can be seen by choosing $A = 0$, $m = 0$ in (1) and rescaling $\tau \rightarrow \tau T$, $x_\mu \rightarrow x_\mu \sqrt{T}$. In Fig. 1 we show the n_L distribution in $D = 3$ and 4 for our working ensembles of loops. We observe the excellent agreement with (8) for large L and the expected systematic deviations for small L . In the following we shall use the actually realized n_L (instead of $L^{-D/2}$) to normalize the loop contributions when necessary, to improve on the systematic errors.

Notice that the number of loops at large L is much smaller than at small L . If large loops are physically important this fact may lead to large statistical errors. One improves on this by using a larger number of trials at high L , producing in this way more loops there. Then we must renormalize their contributions according to the number of trials. The loop frequencies are thus:

$$\nu_L = \frac{2n_L}{(2\pi/D)^{D/2} \mathcal{N}_L} \underset{L \gg 1}{=} L^{-D/2}, \quad \mathcal{N}_L = \text{nr. of trials at } L \quad (9)$$

(the factor 2 appears because on the cubic lattices we only have even loop lengths).

3 Special configurations

For a first test of our approach we use special gauge field configurations: a constant magnetic field in full or half space and a constant topological charge density.

3.1 Homogeneous fields

For homogeneous field configurations the continuum path integral can be analytically evaluated to give the well known Euler-Heisenberg-Schwinger Lagrangian (here in Euclidean space-time):

$$\mathcal{L}(c, b) = \frac{1}{(4\pi)^{D/2}} \int_0^\infty \frac{dT}{T} T^{-D/2} e^{-m^2 T} \frac{cT}{\sinh(cT)} \frac{bT}{\sinh(bT)} \quad (10)$$

where

$$\begin{aligned} c^2 &= \frac{1}{2} \left[\vec{E}^2 + \vec{B}^2 - \sqrt{(\vec{E}^2 + \vec{B}^2)^2 - 4(\vec{E} \vec{B})^2} \right] \\ b^2 &= \frac{1}{2} \left[\vec{E}^2 + \vec{B}^2 + \sqrt{(\vec{E}^2 + \vec{B}^2)^2 - 4(\vec{E} \vec{B})^2} \right] \end{aligned} \quad (11)$$

Of course before performing the T integration one has to renormalize, i.e. to subtract the integrand at $T = 0$ for $D = 3$ and to subtract also its T^2 dependence for $D = 4$

$$\frac{x}{\sinh x} \rightarrow \frac{x}{\sinh x} - 1 - \frac{1}{6}x^2 \quad (12)$$

The constant magnetic field can be realized on the lattice as

$$U_{n,\hat{1}} = \exp \left(2\pi i n_y \frac{k_y}{\hat{N}_y} \right), \quad n_y = 1, \dots, \hat{N}_y = N_y, \quad (13)$$

$$U_{n,\hat{2}} = \exp \left(-2\pi i n_x \frac{k_x}{\hat{N}_x} \right), \quad n_x = 1, \dots, \hat{N}_x = N_x, \quad (14)$$

all other links being 1. This provides a field

$$b_x = 2\pi k_x / \hat{N}_x, \quad b_y = 2\pi k_y / \hat{N}_y, \quad b = b_x + b_y, \quad (15)$$

where k_x, k_y are integers, as required by the periodic boundary conditions.

In Figs. 2 and 3 (left hand plot) we show the quantity:

$$F(L, b) = \frac{1}{n_L} \sum_{i=1}^{n_L} \text{Tr} \left(\prod_{l \in \omega_L^i} U_l \right) \quad (16)$$

for various lattices. From (10,11) (with $B_z = b$, all other fields zero) this should be

$$F_c(L; b) = \frac{u}{\sinh u}, \quad u = \frac{bL}{2D}. \quad (17)$$

For small fields the points fall well together and reproduce the continuum result (17). They also scale well with the lattice size - which, by (13-15), fixes the roughness of the potential, hence indirectly the lattice spacing. For large fields deviations are seen on small lattices, which should be explained by lattice artifacts and signal the departure from the continuum. Notice that the data in the figures are plotted against bL , therefore for large b we see the small L contributions.




Figure 2: Contributions of loops of given length, $F(L, b)$ vs $x \equiv bL$, in a homogeneous magnetic field b for various b in units of $\pi/8$ on a 32^3 lattice in $D = 3$ (left plot) and in units of $\pi/6$ on a 24^4 lattice in $D = 4$ (right plot).

We also consider SU(2) configurations of the form:

$$U_{n,\hat{1}} = \exp\left(2\pi i \sigma_3 n_y \frac{k_y}{N_y}\right), \quad n_y = 1, \dots, N_y, \quad (18)$$

$$U_{n,\hat{3}} = \exp\left(-2\pi i \sigma_3 n_t \frac{k_t}{N_t}\right), \quad n_t = 1, \dots, N_t, \quad (19)$$

all other links being 1. We take $N_\mu = N$ and

$$k_y = k_t = k; \quad b = 2\pi k/N \quad (20)$$

where k is an integer. Such a configuration has $\vec{E} = \vec{B} = (0, 0, b)$ and hence topological charge density

$$q = \frac{2k^2}{N^2} = \frac{b^2}{8\pi^2} \quad (21)$$

homogeneously distributed over the lattice. We calculate

$$F(L, b) = \frac{1}{n_L} \sum_{i=1}^{n_L} \text{Tr} \left(\prod_{l \in \Gamma_i^L} U_l \right), \quad (22)$$

shown in Fig. 3, right hand plot. From (10,11) this should be

$$F_c(L, b) = \frac{u^2}{\sinh^2 u}, \quad u = \frac{bL}{4D}. \quad (23)$$

Again we see good agreement and scaling both with the field and with the lattice size.




Figure 3: Contributions of loops of given length, $F(L, b)$ vs $x \equiv bL$ for various lattices N^4 in $D = 4$: in a homogeneous magnetic field b (b in units of $4\pi/N$) (left plot) and in a field of constant topological charge density $q = \frac{b^2}{8\pi^2}$, b in units of $2\pi/N$ (right plot).

3.2 Magnetic field in a half space

Let us consider now a magnetic field in $D = 3$ in a half space (x-direction) pointing into the z -direction. Take $\hat{N}_x = N_x/2$ in (14) and $m = 0$. We calculate

$$\mathcal{L}(b, x) = \sum_L \frac{1}{L} \nu_L F(L, b, x); \quad F(L, b, x) = \frac{1}{n_L} \sum_{i=1}^{n_L} \text{Tr} \left(\prod_{l \in \tilde{\omega}_L^i(x)} U_l \right) \quad (24)$$

where ν_L are the correct loop frequencies (9). Here the loops in $\tilde{\omega}_L^i(1)$ are obtained from the original ensemble $\omega_L^i(1)$ by approximate centering at the point $x_\mu = 1$ (and are hence confined in a radius $\sim \sqrt{L}$ around this point). Then the $\tilde{\omega}_L^i(x)$ are obtained from the $\tilde{\omega}_L^i(1)$ by translation with x lattice units in the x -direction. We denote

$$\mathcal{L}(L_1, L_2, b, m, x) = \sum_{L=L_1}^{L_2} \frac{1}{L} \nu_L F(L, b, m, x) \quad (25)$$

From the original T - integral (1), as discretized in (2), we should have:

$$\mathcal{L} \left(\frac{\lambda_1}{b}, \frac{\lambda_2}{b}, b, \mu\sqrt{b}, \xi\sqrt{b} + x_0 \right) = b^{-\frac{3}{2}} f(\lambda_1, \lambda_2, \mu, \xi). \quad (26)$$

where x_0 ($\xi = 0$) is the separation between the regions with and without magnetic field. We illustrate in Fig. 4 the test of this scaling relation. We find out that it is rather well fulfilled if we take $x_0 = \frac{N_x}{2} + 0.5$ – notice, that according to (14,15) the field extends with half strength in the region $\frac{N_x}{2}, \frac{N_x}{2} + 1$ before vanishing.

On physical grounds, see also [11], we expect

$$f(0, \infty, \mu, \xi) \propto e^{-M(\mu)\xi} \quad (27)$$




Figure 4: *Scaling propagator $f(\lambda_1, \lambda_2, \mu, \xi)$ eq. (26) for $\mu = 0$, various b (in units of $\pi/16$). Left plot: f vs ξ for various summation intervals $\lambda_1 - \lambda_2$. Right plot: f vs λ_1 for $\mu = 0$, $\lambda_2 = 180$ and $\xi = -5$ (non-vanishing field region).*

In the actual calculation we have $0 < \lambda_{1,2} < \infty$ and discretization artifacts, which bring systematic errors on M . On Fig. 4, right hand plot, we illustrate the dependence of f on λ_1 ; M itself, however, appears less sensitive – see Fig. 4, left hand plot, and Fig. 5. In Fig. 5 we show the “effective mass”

$$M_{\text{eff}}(\lambda_1, \lambda_2, \mu, \xi) = b^{-1/2} \ln \frac{\mathcal{L}\left(\frac{\lambda_1}{b}, \frac{\lambda_2}{b}, b, \mu\sqrt{b}, x\right)}{\mathcal{L}\left(\frac{\lambda_1}{b}, \frac{\lambda_2}{b}, b, \mu\sqrt{b}, x+1\right)} \quad (28)$$

We observe reasonably good scaling, at least for not too large b and m (where small loops dominate and therefore the discretization artifacts are enhanced). Notice that the $m > 0$ results continue nicely to the $m = 0$ ones, indicating that we do not need to take extrapolations but can directly work at $m = 0$. We read off from these figures $M(0) = 1.7 \pm 0.2$. The error is estimated from the sensitivity to the other parameters. Since we expect statistical errors to be smaller than this we did not attempt a jack knife analysis. Note that our result for $M(0)$ is about a factor 2 larger than that of ref. [11]. This disagreement is beyond the estimated error.

To understand the roots of the systematic error we observe that in order to perform the T integral, we need reliable estimates over the full T range. Consider the simpler case of a constant field in the full space: For small T below $T = \frac{2}{D}$ (or, respectively, $T = \frac{3}{2D}$ in the spirit of a Riemann sum) corresponding to our smallest path with $L = 4$ the integrand factor $(bT/\sinh bT - 1)$ can be safely approximated by $-\frac{1}{6}(bT)^2$ for $bT < 1$. For large paths with $L \geq L_{Max} = 10D/b$ corresponding to $bT \geq 5$ the $bT/\sinh bT$ can be neglected against -1 in the integrand, i.e. we only need paths in the outer field below L_{Max} . The integrand above $T_{Max} = L_{Max}/2D$ is easily evaluated (as well as the corresponding L sum producing a ζ function). Both estimates indicate that the integrand at small and large L can be approximated by extrapolation.




Figure 5: *Effective mass $M_{\text{eff}}(\lambda_1, \lambda_2, \mu, \xi)$ eq. (28) for various summation intervals $\lambda_1 - \lambda_2$ and various b (in units of $\pi/16$): vs ξ for $\mu = 0$ (left plot) and vs μ^2 , $\xi \simeq 0.5$ (right plot).*

4 Discussion and Conclusions

We have introduced the Random Walk Worldline method as a statistical summation of the loop expansions for (logarithms of) determinants, and have given examples for effective actions of rather simple gauge fields. A more exciting application would be to lattice discretized nontrivial field configurations like instantons, sphalerons, bounce solutions.

We had a scalar particle in the loop but it is very simple to change to fermions and gauge bosons, just adding a spin term in the worldline Lagrangian containing the outer field [3]. In the case of a (nonabelian) gauge field background spin 1/2 corresponds to a term $i\sigma_{\mu\nu}\mathbf{F}_{\mu\nu}$, spin 1 to $2i\mathbf{F}_{\mu\nu}$ (Feynman gauge). These terms add knots to the Wilson loop and can be discretized as U -plaquettes $U_\mu U_\nu U_\mu^+ U_\nu^+$ to be joined to the path ω_L in all possible ways with a $\sigma_{\mu\nu}$ or 2-factor at the conjunction for spin 1/2 or spin 1, respectively. E.g. the Euler-Heisenberg formula for fermions is thus obtained from the bosonic case very easily. There is an overall Dirac and color trace for fermions and a Lorentz and color trace for gauge bosons in the loop. The latter of course require U -connections in the adjoint representation. The spin term can be further modified by introducing Grassmann variables. This is very useful in analytical calculations, in particular if worldline supersymmetry is used [17], [3]. Also scalar/pseudoscalar and axial vector couplings to spin 1/2 fermions can be treated that way.

Alternatively we can stay with the direct expansion of the fermionic (logarithm of the) determinant and perform the Dirac traces the way it is done for the hopping parameter expansion [14].

Even more interesting is the application of the method in the case of a fluctuating gauge field background in 3- and 4-dimensional QCD and 3-dimensional thermal electroweak theory. This leads to genuine nonperturbative phenomena.

In the general case the quenched matter free energy is given by (5) where we use the ensemble of loops generated by random walk. Varying β we tune the lattice spacing. The

size of the lattice should be chosen such as to accommodate the loops which contribute significantly to (5).

Of course one can discuss strongly interacting gauge theories like QCD fully with lattice methods. Still it is a relevant question how the elementary fermion/gauge field propagators (here in a loop) are deformed in presence of such backgrounds. The answer might allow a semianalytical treatment of some aspects.

In this context the discussion of chiral symmetry breaking seems to be most promising. For our simple case of a constant B -field this is easy to explain in an analytical treatment: calculate e.g. in $d = 3$ with the Banks-Casher formula for the fermion in the loop

$$\begin{aligned} \langle \bar{\psi}\psi \rangle &= -\frac{d}{dm}\Gamma^{(1)}(B)\Big|_{m^2=0} = -2(2m)\int_0^\infty \frac{dT}{T} \frac{1}{(4\pi T)^{3/2}} \frac{BT}{\tanh BT} e^{-m^2 T} \Big|_{m=0} \\ &= -\frac{8mB}{(4\pi)^{3/2}m} \int_0^\infty dx e^{-x^2} = \frac{B}{2\pi} \end{aligned} \quad (29)$$

(this is well known [18]). A discretization in case of a fluctuating gauge field background appears to be very promising [19]. One also may like to develop some modified perturbation theory with an IR cutoff like in the stochastic vacuum model [20] and thus discuss a “magnetic gauge boson mass” [8], perhaps with a tachyonic component, in this context.

Acknowledgment:

We should like to thank Martin Reuter and Werner Wetzel for discussions. The calculations have been done on the VPP300 Computer at the University of Karlsruhe.

References

- [1] R. P. Feynman, Phys. Rev. **80** (1955) 440
- [2] M. J. Strassler, Nucl. Phys. **B385** (1992) 145
- [3] M. G. Schmidt, C. Schubert, Phys. Lett. **B318** (1993) 438;
M. Reuter, M. G. Schmidt, C. Schubert, Ann. of Physics **259** (1997) 319, and refs. therein (hep-th 9610191)
- [4] W. Heisenberg, H. Euler, Z. Phys. **38** (1936) 714; J. Schwinger, Phys. Rev. **82** (1951) 664
- [5] D. Fliegner, M. G. Schmidt, C. Schubert, Z. Phys. **C64** (1994) (hep-ph/9401221);
D. Fliegner, P. Haberl, M. G. Schmidt, C. Schubert, Ann. Phys. **264** (1998) 51 (hep-th/9707189)
- [6] J. Kripfganz, A. Laser, M. G. Schmidt, Nucl. Phys. **B433** (1995) 467; Z. Phys. **C73** (1997) 353 (hep-ph/9512340)
- [7] J. Baacke, Phys. Rev. **D52** (1995) 6760 (hep-ph/9503350)
- [8] S. J. Huber, M. Reuter, M. G. Schmidt, Phys. Lett. **B462** (1999) 158 (hep-ph/9906358)

- [9] M. G. Schmidt, C. Schubert, Phys. Lett. **B 331** (1994) 69 (hep-th/9403158); Phys. Rev. **D53** 91996) 2150 (hep-th/9410100)
- [10] H. T. Sato, M. G. Schmidt, C. Zuhlten, Nucl. Phys. **B579** (2000) 492 (hep-th/00 030 70)
- [11] H. Gies, K. Langfeld, Nucl. Phys. **B613** (2001) 353 (hep-ph/0102185)
- [12] C. Itzykson, J. M. Drouffe, Statistical Field Theory (Cambridge University Press)
- [13] N. Kawamoto, Nucl. Phys. **B90**[FS3] (1981) 617
- [14] I.-O. Stamatescu, Phys. Rev. **D25** (1982) 1130
- [15] T. Bakeyev and Ph. De Forcrand, hep-lat/0008006
- [16] F. Karsch, E. Seiler, I.-O. Stamatescu, Nucl. Phys. **B271** (1986) 349
- [17] L. Brink, S. Deser, B. Zumino, P. DiVecchia, P. Howe, Phys. Lett. **B64B** (1976) 435;
L. Brink, P. DiVecchia, P. Howe, Nucl. Phys. **B118** (1977) 76
- [18] V. P. Gusynin, hep-th/0001070;
V. P. Gusynin, V. A. Miransky, I. A. Shovkovy, Nucl. Phys. **B462** (1996) 249
- [19] M. Reuter, M. G. Schmidt, I.-O. Stamatescu, W. Wetzel, work in progress.
- [20] H. G. Dosch, Prog. Part. Nucl. Phys. **33** (1994) 121;
Y. A. Simonov, Schlading lecture 1996 and further references therein